



تقرير نتائج تحاليل د.جيناتكس
للمية والرجيم الغذائي:



كيف تقرأ وتستفيد من المعلومات التي يتضمنها تقرير نتائج تحليل

الحمية والرجيم الغذائي:



صحة الإنسان تعتمد على عمليات معقدة جداً ترتبط ارتباطاً وثيقاً بين سماته الجينية والعوامل البيئية المتغيرة (نمط الحياة، البرنامج الغذائي، الرياضة والحركة، الضغط النفسي، الإرهاق والإجهاد، إلخ).

يجب معرفة أن هذه المقدمة هي خاصة بك، لذلك تمت كتابة ملخص التقرير بلغة سهلة يستطيع الفرد الغير متخصص طبياً في فهمها واستيعاب محتواها. ولكن يجب أن تعلم أيضاً بأن باقي التقرير مقدم للأخصائي الصحي، والذي بدوره يستطيع استخلاص محتويات التقرير على شكل برنامج صحي مخصص لك أنت فقط، اعتماداً على تحليل سماتك الجينية التي قام مركز بندرجين الطبي باستكشافها لك.

١) كلمة "محمتمل" تم استخدامها بكثرة في هذا التقرير. ماذا تعني؟

يعلم كثير من الناس أن ارتفاع معدل الكوليسترول مثلاً يرتبط بالإصابة بأمراض القلب وانسداد الشرايين، ومع ذلك نجد أن بعض الناس ممن يعانون من ارتفاع الكوليسترول ولكنهم لا يعانون من أي مشاكل قلبية أو وعائية. أيضاً، التدخين يؤدي إلى سرطان الرئة، ولكن ليس مع الجميع. لهذا، يجب أن نعلم أن بعض السمات الجينية لدى الإنسان تؤدي غالباً إلى صفات معينة، ولكن ليس دائماً. لذلك كلمة "محمتمل" في هذا التقرير تعني أن السمة الجينية ستؤدي غالباً إلى النتيجة المتوقعة، ولكن يمكن لبعض العوامل المختلفة تغيير ذلك.

٢) ماذا تعني كلمة "متوسط" في هذا التقرير؟

متوسط تعني ليس مرتفعاً وليس منخفضاً، ولكن في منطقة تقارب المنتصف نسبياً، مثلاً: متوسط احتمالية اكتساب الوزن. بالنسبة للسمات الجينية - خصوصاً بالمقارنة مع النتائج العلمية لشريحة كبيرة من الناس - كلمة متوسط تعني طبيعي أو نمطي.

٣) كيف أعرف ما هي السمات الوراثية الخاص بي أنا لكل بند من بنود التحليل الذي قمت به؟

لقد قمنا بوضع علامة (✓) أمام التفسير الذي ينطبق على حالتك. وطبعاً قمنا بعرض بقية الاحتمالات حتى يمكنك معرفة ماهي النتائج المتوقعة لكل بند، وكذلك لتتمكن من مقارنة نتائجك الشخصية مع النتائج "المتوسطة" علمياً.

٤) كيف توصلتم لهذه النتائج؟

لقد قمنا بتحليل الحمض النووي الخاص بك وربط نتائج المتغيرات (الطفرات) الجينية الموجودة لك مع نتائج أبحاث علمية منشورة في قواعد البيانات الطبية العالمية المتخصصة في تحليل سلسلة الحمض النووي (DNA). يمكنك الاطلاع على قائمة بالمراجع العلمية الموجودة على موقعنا الإلكتروني للتعرف أكثر عن الدراسات العلمية والطبية في مجال الفحص الجيني والوراثي.

٥) بعض الجينات تكون ذات استدلال مفيد والبعض الآخر ذات استدلال غير مفيد لنفس الصفة. كيف يمكن تفسير ذلك؟

يحتوي الحمض النووي للإنسان على حوالي ٢٠-٣٠ ألف جين. معظم الخصائص والوظائف العضوية والحيوية في جسم الإنسان يتم تنظيمها عن طريق مجموعة من الجينات (وليس جين واحد فقط لكل وظيفة). لذلك كانت بعض السمات الجينية ذات استدلال مفيد لصفة معينة، وجينات أخرى ذات استدلال غير مفيد لذات الصفة. أيضاً يجب أن تعلم أن ليس كل جين يقوم بالتأثير على الصفة بشكل قياسي، فبعض السمات الجينية تؤثر بشكل كبير وبعض السمات الجينية تؤثر بشكل أقل.

٦) بعض العبارات في التقرير تم تلويها بالأحمر وبعضها تم تلويها بالأخضر، لماذا؟

السمات الجينية التي يعتبر لها استدلال مفيد وإيجابي فيما يتعلق بالفائدة الغذائية الخاصة بك قد تم تعليمها بالأخضر. وكذلك السمات الجينية التي يعتبر لها استدلال غير مفيد وسلبي فيما يتعلق بالفائدة الغذائية الخاصة بك قد تم تعليمها بالأحمر. باقي الاستدلالات التي يتم ربطها علمياً (إما إيجاباً أو سلباً) تم تعليمها بالأسود.

نتائج فحص السمات الوراثية الخاصة بكم للحمية والرجيم الغذائي:

| النتائج المحتملة | نتيجتك | السمة الجينية |
|---|--------|--|
| السمات المتعلقة بالحفاظ على الوزن | | |
| متوسط احتمالية الرغبة في الأكل الزائد واكتساب الوزن | ✓ | قابلية الزيادة في الوزن: قابلية زيادة الوزن نتيجة تخزين كميات كبيرة من الطاقة الخلوية الناتجة من أيض المواد الغذائية المختلفة. |
| انخفاض احتمالية الرغبة في الأكل الزائد واكتساب الوزن | | |
| ارتفاع احتمالية الرغبة في الأكل الزائد واكتساب الوزن | | |
| متوسط احتمالية الزيادة في الأكل | ✓ | قابلية الزيادة في الأكل: قابلية الإحساس المستمر بالنهم أو الرغبة في الاستمرار في الأكل حتى بعد الوصول الطبيعي لحالة الشبع. |
| انخفاض احتمالية الزيادة في الأكل | | |
| ارتفاع احتمالية الزيادة في الأكل | | |
| متوسط احتمالية استهلاك الأطعمة الدسمة والدهنية | ✓ | قابلية تناول الأطعمة الدهنية: بعض الأشخاص الذين يحملون سمات جينية محددة لديهم ميول طبيعي للأطعمة الدسمة والدهنية ويفضلونها على الأطعمة الصحية أو المنزوعة-المخففة من الدسم، مما يؤدي إلى اكتساب الوزن. |
| انخفاض احتمالية استهلاك الأطعمة الدسمة والدهنية | | |
| ارتفاع احتمالية استهلاك الأطعمة الدسمة والدهنية | | |
| متوسط احتمالية استهلاك الأطعمة السكرية | ✓ | قابلية تناول السكريات: بعض الأشخاص الذين يحملون سمات جينية محددة لا يقومون بالتحسس من الطعم السكري المرتفع، لذلك نجد أنهم يتناولون الأطعمة والمواد النشوية والسكرية أكثر من غيرهم مما يؤدي إلى اكتساب الوزن. |
| انخفاض احتمالية استهلاك الأطعمة السكرية | | |
| ارتفاع احتمالية استهلاك الأطعمة السكرية | | |
| متوسط احتمالية الرغبة في تناول الخضروات | ✓ | قابلية تناول الخضروات: بعض الأشخاص الذين يحملون سمات جينية محددة لا يحبون تناول الخضروات بسبب الطعم المر أو المعدني الموجود في العديد من الخضروات، على الرغم من أن تناول الخضروات يحافظ على الوزن المثالي. |
| انخفاض احتمالية الرغبة في تناول الخضروات | | |
| ارتفاع احتمالية الرغبة في تناول الخضروات | | |
| متوسط احتمالية اكتساب الوزن نتيجة تناول النشويات | ✓ | تناول الأطعمة السكرية واكتساب الوزن: بعض الأشخاص الذين يحملون سمات جينية محددة يكتسبون الوزن بسرعة نتيجة تناول السكريات. |
| انخفاض احتمالية اكتساب الوزن نتيجة تناول النشويات | | |
| ارتفاع احتمالية اكتساب الوزن نتيجة تناول النشويات | | |
| متوسط احتمالية اكتساب الوزن نتيجة تناول الدهون المشبعة | ✓ | تناول الدهون المشبعة واكتساب الوزن: بعض الأشخاص الذين يحملون سمات جينية محددة يكتسبون الوزن بسرعة نتيجة تناول الدهون المشبعة. |
| انخفاض احتمالية اكتساب الوزن نتيجة تناول الدهون المشبعة | | |
| ارتفاع احتمالية اكتساب الوزن نتيجة تناول الدهون المشبعة | | |
| متوسط احتمالية اكتساب الوزن نتيجة تناول MUFA | ✓ | تناول الدهون الأحادية الغير مشبعة (MUFA) واكتساب الوزن: بعض الأشخاص الذين يحملون سمات جينية محددة يكتسبون الوزن بسرعة نتيجة تناول الدهون الأحادية الغير مشبعة. |
| انخفاض احتمالية اكتساب الوزن نتيجة تناول MUFA | | |
| ارتفاع احتمالية اكتساب الوزن نتيجة تناول MUFA | | |
| متوسط احتمالية اكتساب الوزن نتيجة تناول PUFA | ✓ | تناول الدهون المتعددة الغير مشبعة (PUFA) واكتساب الوزن: بعض الأشخاص الذين يحملون سمات جينية محددة يكتسبون الوزن بسرعة نتيجة تناول الدهون الأحادية الغير مشبعة. |
| انخفاض احتمالية اكتساب الوزن نتيجة تناول PUFA | | |
| ارتفاع احتمالية اكتساب الوزن نتيجة تناول PUFA | | |

نتائج فحص السمات الوراثية الخاصة بكم للمحبة والرجيم الغذائي:

| النتائج المحتملة | نتيجتك | السمة الجينية |
|---|--------|---|
| السمات المتعلقة بالحفاظ على الوزن | | |
| متوسط احتمالية خسارة الوزن نتيجة تناول البروتينات | ✓ | تناول البروتينات وخسارة الوزن: بعض الأشخاص الذين يحملون سمات جينية محددة يخسرون الوزن بسرعة نتيجة تناول كمية أكبر من البروتينات. |
| ارتفاع احتمالية خسارة الوزن نتيجة تناول البروتينات | | |
| متوسط احتمالية اكتساب الوزن بسرعة بعد خسارته | ✓ | قابلية إعادة اكتساب الوزن بعد خسارته: بعض الأشخاص الذين يحملون سمات جينية محددة يكتسبون الوزن بسرعة بعد برنامج طويل ومجهد لخسارة الوزن، مقارنة بالبعض الآخر الذين يحافظون على وزنهم الجديد دون زيادة كبيرة لفترات أطول. |
| انخفاض احتمالية اكتساب الوزن بسرعة بعد خسارته | | |
| ارتفاع احتمالية اكتساب الوزن بسرعة بعد خسارته | | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين أ في غذائك | | احتياجك لفيتامين أ: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين أ. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين أ في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين ب١٢ في غذائك | | احتياجك لفيتامين ب١٢: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين ب١٢. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين ب١٢ في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين ب٦ في غذائك | ✓ | احتياجك لفيتامين ب٦: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين ب٦. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين ب٦ في غذائك | | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين ب٩ في غذائك | ✓ | احتياجك لفيتامين ب٩: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين ب٩. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين ب٩ في غذائك | | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين سي في غذائك | ✓ | احتياجك لفيتامين سي: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين سي. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين سي في غذائك | | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين د في غذائك | | احتياجك لفيتامين د: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين د. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين د في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين إي في غذائك | | احتياجك لفيتامين إي: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين إي. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين إي في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية فيتامين ك في غذائك | | احتياجك لفيتامين ك: بعض الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بفيتامين ك. |
| مرتفع: تحتاج إلى زيادة كمية من فيتامين ك في غذائك | ✓ | |
| منخفض: تحتاج إلى كمية أقل من الكالسيوم في غذائك | | احتياجك للكالسيوم: بعض الأشخاص الذين يحملون سمات جينية محددة لا يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالكالسيوم. |
| متوسط: تحتاج إلى كمية قياسية من الكالسيوم في غذائك | | |
| مرتفع: تحتاج إلى زيادة كمية الكالسيوم في غذائك | ✓ | |

نتائج فحص السمات الوراثية الخاصة بكم للحمية والرجيم الغذائي:

| النتائج المحتملة | نتيجتك | السمة الجينية |
|---|--------|--|
| طبيعي: تحتاج إلى كمية قياسية من الكولين في غذائك | | احتياجك لمادة الكولين: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالكولين. |
| مرتفع: تحتاج إلى زيادة كمية الكولين في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية من النحاس في غذائك | | احتياجك لمادة النحاس: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالنحاس. |
| مرتفع: تحتاج إلى زيادة كمية النحاس في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية من الحديد في غذائك | | احتياجك لمادة الحديد: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالحديد. |
| مرتفع: تحتاج إلى زيادة كمية الحديد في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية من المغنيسيوم في غذائك | | احتياجك لمادة المغنيسيوم: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالمغنيسيوم. |
| مرتفع: تحتاج إلى زيادة كمية المغنيسيوم في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية من الفوسفات في غذائك | | احتياجك لمادة الفوسفات: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالفوسفات. |
| مرتفع: تحتاج إلى زيادة كمية الفوسفات في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية من الزنك في غذائك | | احتياجك لمادة الزنك: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالزنك. |
| مرتفع: تحتاج إلى زيادة كمية الزنك في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية من السيلينيوم في غذائك | | احتياجك لمادة السيلينيوم: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بالسيلينيوم. |
| مرتفع: تحتاج إلى زيادة كمية السيلينيوم في غذائك | ✓ | |
| طبيعي: تحتاج إلى كمية قياسية من مضادات الأكسدة | | احتياجك لمضادات الأكسدة: بعض الأشخاص الذين يحملون سمات جينية محددة يحتاجون إلى زيادة استهلاك الأطعمة الغنية بمضادات الأكسدة. |
| مرتفع: تحتاج إلى زيادة كمية مضادات الأكسدة | ✓ | |
| مرتفع: زيادة الرغبة في استهلاك الكافيين | | استهلاك الكافيين: بعض الأشخاص الذين يحملون سمات جينية محددة تزيد لديهم الرغبة في استهلاك الكافيين بكميات مرتفعة. |
| منخفض: انخفاض الرغبة في استهلاك الكافيين | | |
| طبيعي: تحتاج إلى كمية قياسية من الكافيين | ✓ | |

نتائج فحص السمات الوراثية الخاصة بكم للحمية والرجيم الغذائي:

| النتائج المحتملة | نتيجتك | السمة الجينية |
|---|--------|--|
| بطيء: احتمالية التمثيل الغذائي لمادة الكافيين بشكل بطيء | ✓ | تمثيل (أيض) الكافيين: بعض الذين يحملون سمات جينية محددة يقومون بعملية التمثيل الغذائي لمادة الكافيين بشكل بطيء لذلك يعانون من الأرق لفترات أطول بعد تناوله. |
| سريع: احتمالية التمثيل الغذائي لمادة الكافيين بشكل سريع | | |
| مرتفع الحساسية: احتمالية عدم تحمل مركب الجلوتين | ✓ | الحساسية الزائدة لبروتين القمح (الجلوتين): بعض الذين يحملون سمات جينية محددة لا يستطيعون تحمل الأطعمة المحتوية على الجلوتين لذلك يعانون من مضاعفات وتقلصات في الجهاز الهضمي. |
| منخفض الحساسية: احتمالية تحمل مركب الجلوتين | | |
| مرتفع الحساسية: احتمالية عدم تحمل سكر اللاكتوز | ✓ | الحساسية الزائدة لسكر الحليب (اللاكتوز): بعض الذين يحملون سمات جينية محددة لا يستطيعون تكسير اللاكتوز وهضمه بشكل كامل مما يؤدي إلى مضاعفات وتقلصات في الجهاز الهضمي. |
| منخفض الحساسية: احتمالية تحمل سكر اللاكتوز | | |
| انخفاض احتمالية انخفاض ضغط الدم عند تقليل الملح | ✓ | كمية الملح وارتفاع ضغط الدم: بعض الأشخاص الذين يحملون سمات جينية محددة يمكنهم المحافظة على ضغط دم منخفض إذا قاموا بتقليل كمية الملح في برنامجهم الغذائي. |
| متوسط احتمالية انخفاض ضغط الدم عند تقليل الملح | | |
| ارتفاع احتمالية انخفاض ضغط الدم عند تقليل الملح | | |
| منخفض: معدل الاستجابة للريوفلافين منخفضة | ✓ | كمية الريوفلافين وانخفاض ضغط الدم: بعض الذين يحملون سمات جينية محددة يمكنهم المحافظة على ضغط دم منخفض إذا قاموا بزيادة كمية الريوفلافين في برنامجهم الغذائي. |
| مرتفعة: معدل الاستجابة للريوفلافين مرتفعة | | |

INTRODUCTION

Understanding your report

Have you ever wondered why certain people lose or gain more weight compared to others? And why some foods cause uneasiness in some people? Do you want to know which diet suits you best? The answer lies in your genes.

The way we fuel our bodies with the foods we eat are all impacted by our genetic make-up. The old adage “you are what you eat” plays a major role in determining our health and well-being. Food and its nutrients directly and indirectly influence our gene expressions. Genetic variations affecting certain metabolic traits in turn dictate dietary means and requirements. For instance, the response to food varies from individual to individual explaining why some people can eat as much as they want and not gain weight. These factors may be attributed to the large role that genes play in influencing eating behaviours and metabolism of different foods.

Some interesting facts about genes and nutrition:

- Craig Maclean, the famous track cyclist and Olympics Gold medalist, and Novak Djokovic, the famous tennis player, were both diagnosed with Celiac disease (gluten intolerance) and owe their success to a gluten-free diet.
- A study on weight management conducted by Stanford University found that people who eat and exercise according to their genetic predisposition tend to lose two-and-a-half times as much weight as compared to those who do not.
- Approximately 74% of Native American, 90% of Asian Americans, 70% of African Americans and 53% of Mexican Americans are lactose intolerant. Research studies have also showed that there is a considerable reduction in lactase activity among people whose ancestry is from Greek, Italian, Arab, Asian, African, Hispanic or Jewish origin.
- According to Centres for Disease Control and Prevention (CDC), 43% of children and 38% of pregnant women suffer from iron deficiency, 1 in 6 women suffer from Vitamin A deficiency and 17.3% of the global population has Zinc deficiency.
- Fast metabolizers of caffeine, who drank up to 3 cups of coffee per day, have almost 52% lower risk of heart attack as compared to slow metabolizers.

In this report, we profile genes that have been shown to influence nutritional traits like diet and weight management, micronutrient requirements, food intolerance and several other attributes relevant to nutritional well-being.

We hope that this report will help you understand your body better and to align your diet to your genetic type to get the best results.

INTRODUCTION

Understanding your report

Human health is a complex interplay between genetics and the environment (lifestyle, diet, activity, stress, etc.). Your genes, training and diet, all play a vital role in your well-being.

This report is presented in a user friendly language and format. The following tips will help you get the best information value out of the report.


1. The word "likely" is used often in the report. What does it mean?

People generally know that high cholesterol can lead to heart conditions. However, there are individuals with high cholesterol who do not develop heart disease. Similarly, smoking can lead to lung disease, but not always. Hence, certain genetic parameters can lead to certain outcomes but other factors may modify the outcome. "Likely" means, it is more likely that one will see the outcome, but other factors may modify it.

2. What does the term "average" mean in the report?

Average implies neither high nor low, rather an intermediate outcome. For example, average likelihood of weight gain is an intermediate level between high and low likelihood. Average can also be understood in the context of "Normal" or "Typical" or "Moderate"

3. How do I know which result is applicable to me?

Only results with a check mark  are applicable to you, the others are not applicable. All possible outcomes are provided in the table to provide a context to your outcome.

4. Where did the information contained in the report come from?

The genetic markers that are used in this report are based on scientific studies published in international journals. A list of references is available for you to read on our web blog.

5. Some sentences are colored in green and others in red, why?


Attributes that are advantageous in nutritional well-being are indicated in green and those that are not advantageous are in red. Moderate or Neutral outcomes are indicated in black.

6. In the vitamins and Micro Nutrient section, what does normal intake indicate?

Normal intake refers to the Recommended Dietary Allowance (RDA) of the specific vitamin or mineral.

SUMMARY RESULTS

Your diet plan



Key takeaways

| | |
|----------------------------|---|
| Gluten Sensitivity | Gluten Sensitive: Increased risk of being gluten sensitive |
| Lactose Intolerance | Intolerant: Likely to be lactose intolerant |
| Alcohol Flush | Low: Less likely to experience alcohol flush |
| Increase Intake of | Vitamin A, Vitamin E, Vitamin K, Calcium, Choline, Copper, Iron |

Nutrition Results

| TRAIT NAME | YOUR RESULTS | POSSIBLE OUTCOMES |
|---|--------------|--|
| Tendency To Gain Weight A genetic tendency for higher weight gain due to increased energy storage | ✓ | Moderate: Moderately likely to gain weight |
| | | Low: Less likely to gain weight |
| | | High: Highly likely to gain weight |
| Tendency To Overeat A genetic tendency to over consume foods due to increased craving | ✓ | Moderate: Moderately likely to overeat |
| | | Low: Less likely to overeat |
| | | High: Highly likely to overeat |
| Tendency To Prefer Fatty Foods A genetic tendency for lower fat taste perception and overconsumption of fatty foods | | Moderate: Likely to overconsume high fat foods |
| | ✓ | Low: Less likely to overconsume high fat foods |
| | | High: Highly likely to overconsume high fat foods |
| Tendency To Prefer Sweet Foods A genetic tendency for lower sweet taste perception and overconsumption of sweet foods | ✓ | Moderate: Likely to overconsume sweet foods |
| | | Low: Less likely to overconsume sweet foods |
| | | High: Highly likely to overconsume sweet foods |
| Tendency To Prefer Bitter Foods A genetic tendency for high bitter taste perception and low intake of bitter vegetables | ✓ | Moderate: Normal bitter vegetable preference |
| | | Low: Less likely to prefer bitter vegetables |
| | | High: Likely to prefer bitter vegetables |
| Carbohydrate Intake And Weight Gain Tendency A genetic tendency for increased weight gain upon higher carb intake | | Moderate: Likely to gain weight on high carb intake |
| | ✓ | Low: Less likely to gain weight on high carb |
| | | High: Highly likely to gain weight on high carb intake |
| Saturated Fats Intake And Weight Gain Tendency A genetic tendency for higher weight gain upon higher sat fat intake | ✓ | Moderate: Likely to gain weight on high SFA intake |
| | | Low: Less likely to gain weight on high SFA intake |
| | | High: Highly likely to gain weight on high SFA intake |
| Mono Unsaturated Fats Intake And Weight Gain Tendency A genetic tendency for weight gain upon higher MUFA intake | | Moderate: Likely to gain weight with high MUFA intake |
| | | Low: Less likely to gain weight with high MUFA intake |
| | ✓ | High: Highly likely to gain weight with high MUFA intake |
| Poly Unsaturated Fats Intake And Weight Gain Tendency A genetic tendency for weight gain upon higher PUFA intake | | Moderate: Likely to gain weight with high PUFA intake |
| | ✓ | Low: Less likely to gain weight with high PUFA intake |
| | | High: Highly likely to gain weight with high PUFA intake |

| TRAIT NAME | YOUR RESULTS | POSSIBLE OUTCOMES |
|---|--------------|---|
| Protein Intake And Weight Loss Tendency A genetic tendency for increased weight loss upon higher protein intake | ✔ | Moderate: Likely to lose weight on high protein intake |
| | | High: Highly likely to lose weight on high protein intake |
| Tendency To Regain Weight A genetic tendency for rapid weight regain after a weight loss program | ✔ | Moderate: Likely to regain weight after weight loss |
| | | Low: Less likely to regain weight after weight loss |
| | | High: Highly likely to regain weight after weight loss |
| Vitamin A Needs A genetic tendency to require more Vitamin A due to inefficient metabolism of vitamin A | | Need more: Moderately increase vitamin A intake |
| | | Normal: Maintain normal Vitamin A intake |
| | ✔ | Need more: Significantly increase vitamin A intake |
| Vitamin B12 Needs A genetic tendency to require more Vitamin B12 due to inefficient metabolism of vitamin B12 | ✔ | Need more: Moderately increase vitamin B12 intake |
| | | Normal: Maintain normal Vitamin B12 intake |
| | | Need more: Significantly increase vitamin B12 intake |
| Vitamin B6 Needs A genetic tendency to require more Vitamin B6 due to inefficient metabolism of vitamin B6 | | Need more: Moderately increase vitamin B6 intake |
| | ✔ | Normal: Maintain normal Vitamin B6 intake |
| | | Need more: Significantly increase vitamin B6 intake |
| Vitamin B9 Needs A genetic tendency to require more Vitamin B9 due to inefficient metabolism of vitamin B9 | | Need more: Moderately increase vitamin B9 intake |
| | ✔ | Normal: Maintain normal Vitamin B9 intake |
| | | Need more: Significantly increase vitamin B9 intake |
| Vitamin C Needs A genetic tendency to require more Vitamin C due to inefficient metabolism of vitamin C | | Need more: Moderately increase vitamin C intake |
| | ✔ | Normal: Maintain normal Vitamin C intake |
| | | Need more: Significantly increase vitamin C intake |
| Vitamin D Needs A genetic tendency to require more Vitamin D due to inefficient metabolism of vitamin D | ✔ | Need more: Moderately increase vitamin D intake |
| | | Normal: Maintain normal Vitamin D intake |
| | | Need more: Significantly increase vitamin D intake |
| Vitamin E Needs A genetic tendency to require more Vitamin E due to inefficient metabolism of vitamin E | | Need more: Moderately increase vitamin E intake |
| | | Normal: Maintain normal Vitamin E intake |
| | ✔ | Need more: Significantly increase vitamin E intake |
| Vitamin K Needs A genetic tendency to require more Vitamin K due to inefficient metabolism of vitamin K | | Need more: Moderately increase vitamin K intake |
| | | Normal: Maintain normal Vitamin K intake |
| | ✔ | Need more: Significantly increase vitamin K intake |

| TRAIT NAME | YOUR RESULTS | POSSIBLE OUTCOMES |
|---|--------------|---|
| Calcium Needs A genetic tendency to need more or less of Calcium due to differences in metabolism of Calcium | | Need less: You may have higher calcium levels |
| | | Normal: Maintain normal Calcium intake |
| | ✓ | Need more: Significantly increase Calcium intake |
| Choline Needs A genetic tendency to require more Choline due to inefficient metabolism of Choline | | Need more: Moderately increase Choline intake |
| | | Normal: Maintain normal Choline intake |
| | ✓ | Need more: Significantly increase Choline intake. |
| Copper Needs A genetic tendency to require more Copper due to inefficient metabolism of Copper | | Need more: Moderately increase Copper intake |
| | | Normal: Maintain normal Copper intake |
| | ✓ | Need more: Significantly increase Copper intake |
| Iron Needs A genetic tendency to require more Iron due to inefficient metabolism of Iron | | Need more: Moderately increase Iron intake |
| | | Normal: Maintain normal Iron intake |
| | ✓ | Need more: Significantly increase Iron intake |
| Magnesium Needs A genetic tendency to require more Magnesium due to inefficient metabolism of Magnesium | ✓ | Need more: Moderately increase Magnesium intake |
| | | Normal: Maintain normal Magnesium intake |
| | | Need more: Significantly increase Magnesium intake |
| Phosphate Needs A genetic tendency to require more Phosphate due to inefficient metabolism of Phosphate | ✓ | Need more: Moderately increase Phosphate intake |
| | | Normal: Maintain normal Phosphate intake |
| | | Need more: Significantly increase Phosphate intake |
| Zinc Needs A genetic tendency to require more Zinc due to inefficient metabolism of Zinc | ✓ | Need more: Moderately increase Zinc intake |
| | | Normal: Maintain normal Zinc intake |
| | | Need more: Significantly increase Zinc intake |
| Selenium Needs A genetic tendency to require more Selenium due to inefficient metabolism of Selenium | | Need more: Moderately increase Selenium intake |
| | ✓ | Normal: Maintain normal Selenium intake |
| | | Need more: Significantly increase Selenium intake |
| Antioxidant Needs A genetic tendency to require more Antioxidants due to inefficient metabolism of Antioxidants | ✓ | Need more: Moderately increase antioxidants intake |
| | | Normal: Maintain normal antioxidants intake |
| | | Need more: Significantly increase antioxidants intake |
| Caffeine Consumption A genetic tendency for variation in caffeine consumption due to difference in preference | | High: Likely to consume more caffeine |
| | | Low: Likely to consume less caffeine |
| | ✓ | Moderate: Likely to consume moderate amount of caffeine |

| TRAIT NAME | YOUR RESULTS | POSSIBLE OUTCOMES |
|---|--------------|---|
| Caffeine Metabolism A genetic tendency for variation in caffeine clearance due to varied metabolism | ✔ | Slow: Likely to be a slow metabolizer of caffeine |
| | | Fast: Likely to be a fast metabolizer of caffeine |
| Gluten Sensitivity A genetic tendency for variation in response to gluten due to varied sensitivity | | Gluten insensitive: Unlikely to be gluten sensitive |
| | ✔ | Gluten Sensitive: Increased risk of being gluten sensitive |
| Lactose Intolerance A genetic tendency for difference in response to lactose due to varied tolerance | | Tolerant: Less likely to be lactose intolerant |
| | ✔ | Intolerant: Likely to be lactose intolerant |
| Salt Intake And Blood Pressure Sensitivity A genetic tendency for lower blood pressure due to salt sensitivity | ✔ | Moderate: Likely to have lower BP on low salt diet |
| | | Normal: Likely to slightly reduce BP on low salt diet |
| | | High: Highly likely to have lower BP on low salt diet |
| Riboflavin And Blood Pressure Response A genetic tendency for lower blood pressure on increased riboflavin intake | ✔ | Insensitive: Less likely to have lower BP on high Riboflavin intake |
| | | Sensitive: Highly likely to have lower BP with high Riboflavin Intake |
| Alcohol Flush A genetic tendency for variation in response to alcohol intake due to reduced clearance of acetaldehyde | ✔ | Low: Less likely to experience alcohol flush |
| | | Moderate: Moderately likely to experience alcohol flush |
| | | High: Highly likely to experience alcohol flush. |

TENDENCY TO GAIN WEIGHT

People of certain genetic type have more of a tendency to gain weight than others due to their genetic makeup. Genes that regulate metabolic rate, energy expenditure and energy storage influence the tendency to store calories versus expending calories. Certain genes that once offered a survival advantage to our ancestors by storing calories for leaner times have now become liabilities in the age of surplus food availability. Your genetic profile influences the tendency of your body to store or expend more calories.



Moderate: Moderately likely to gain weight

| Genes tested | Recommendations |
|---|--|
| <p><i>MAF, UCP2, TMEM18, SH2B1, SEC16B, ADIPOQ, ADRB1, ADRB21, LEP, ETV5, GNPDA2, PCSK1, NEGR1, MTCH2, ADRB2, KCTD15, BDNF, MC4R, FTO, UCP3</i></p> | <ul style="list-style-type: none"> • You have a moderate genetic tendency to gain weight. • Ensure a healthy balance between energy intake and physical activity to maintain a healthy weight. |

TENDENCY TO OVEREAT

People of certain genetic type have more of a tendency to over consume foods. We analyze genes that are known to influence various hunger and satiety hormones such as Leptin, Ghrelin and Neuropeptides, which influence neurological aspects of feeding, producing effects such as persistent hunger, excessive snacking, preference for high calorific food and emotional eating.



Moderate: Moderately likely to overeat

| Genes tested | Recommendations |
|---|---|
| <p><i>CLOCK, MC4R, FTO, TAS2R38, DRD2</i></p> | <ul style="list-style-type: none"> • You may have a moderate genetic tendency to overeat. • Practice mindful eating and eat high fibre or high protein snacks to increase feeling of fullness |

TENDENCY TO PREFER FATTY FOODS

People of certain genetic type tend to over consume fatty foods due to a lower ability to perceive fats. In studies, people with lower fat perception ability were found to rate the fat content of food consistently lower than the actual fat content. Eating high quantities of fatty food can lead to weight gain and other health conditions.



Low: Less likely to overconsume high fat foods

| Genes tested | Recommendations |
|---------------------------|---|
| <p><i>CD361, CD36</i></p> | <ul style="list-style-type: none"> • You may have an increased genetic ability to taste fats and are less likely to over consume high fat foods. • You may have a natural genetic preference for low fat foods and increased intake could be due to other factors like access to high fat food. • Consciously reduce intake of fried and oily foods. • Look for 0 g trans fat on the Nutrition Facts label and no hydrogenated oils in the ingredients list |

TENDENCY TO PREFER SWEET FOODS

People of certain genetic type tend to over consume sweet foods due to low sensitivity to sweet taste, either due to lesser number or reduced sensitivity of sweet taste receptors on their tongue. Sugary foods are rich in calories and can cause insulin resistance, leading to weight gain and other health conditions.



Moderate: Likely to overconsume sweet foods

| Genes tested | Recommendations |
|--|---|
| <p><i>TAS2R381, TAS1R31, TAS1R3, TAS1R21, TAS1R2, GLUT2, TAS2R382, TAS2R38</i></p> | <ul style="list-style-type: none"> • You may have a slightly decreased genetic sweet taste sensitivity and are likely to consume more sweet foods. • Consciously reduce intake of sweets. • High sugar intake increases risk for obesity and diabetes. • Snack on dry fruits, fruits and green leafy vegetables to reduce sugar cravings. • Drink plenty of water. |

TENDENCY TO PREFER BITTER FOODS

Bitter taste perception is due to genetic variations in taste receptors. When food is chewed in the mouth, molecules such as phenylthiocarbamide bind to the taste receptors present in the tongue, which gives rise to the bitter taste. People of certain genetic type tend to avoid bitter vegetables due to higher sensitivity to bitter taste. Vegetables are low calorie and high fibre foods, which help in weight management and have several health benefits.



Moderate: Normal bitter vegetable preference

| Genes tested | Recommendations |
|---|--|
| <p><i>TAS2R38, TAS2R381, TAS2R382</i></p> | <ul style="list-style-type: none"> • You may have a genetic tendency to prefer bitter vegetables which could be due to moderate sensitivity to bitter taste. • Add natural sweeteners, lemon juice or spices to flavor bitter tasting green leafy vegetables, be creative with vegetable preparations. • Bitter foods help absorb nutrients, reduce sugar cravings and balance appetite. • Eat vegetables and fruits of different colors |

CARBOHYDRATE INTAKE AND WEIGHT GAIN TENDENCY

Carbohydrates are the main sources of energy and they provide the kilocalories for weight maintenance. 45-65% of total calories are the recommended intake of carbohydrates with starch and sugars being the major carbohydrates. Corn, rice, potatoes, pasta and breads are sources of starch. Fruits and fruit juices have natural sugars while desserts, candies and soft drinks have added sugar. Carbs are considered as weight increasing foods, but that's not true for everyone. People of certain genetic type tend to gain more weight upon consuming carbohydrate rich foods than others. These individuals can better maintain weight by reducing the amount of carbs in their diet. Eat a balanced diet. Choose complex carbohydrates such as fruits, vegetables, legumes and whole grain.



Low: Less likely to gain weight on high carb

| Genes tested | Recommendations |
|--|--|
| <p><i>FTO, TCF7L2, SEC16B, AMY1, FABP2, FAIM2, FLJ35779, FTO1, RBJ, LRRN6C</i></p> | <ul style="list-style-type: none"> • You may have a low genetic risk for weight gain on a high carbohydrate diet. • Eat a balanced diet. • Choose complex carbohydrates such as fruits, vegetables, legumes and whole grain |

SATURATED FATS INTAKE AND WEIGHT GAIN TENDENCY

Saturated fats are a type of fat that are largely solid at room temperature as they are saturated with hydrogen molecules. Meat and dairy products are rich sources of saturated fats. A high intake of saturated fats is associated with an increase in LDL cholesterol levels in the body. The American Heart Association recommends no more than 5-6% of calories from saturated fat from the daily diet. People of certain genetic type tend to gain more weight upon consuming saturated fat rich foods than others. These individuals can better maintain weight by reducing the amount of saturated fats in their diets



Moderate: Likely to gain weight on high SFA intake

| Genes tested | Recommendations |
|-------------------|--|
| <i>FTO, APOA2</i> | <ul style="list-style-type: none"> • You may have a genetic tendency to have a moderately higher BMI on a high fat diet. • Lowering fat in the diet will help your weight loss goals. • Try to lower consumption of high calorie, processed foods like pizzas, burgers, french-fries, soda, pastries, cookies, candies, cream and cheese sauces; substitute recipes with low-fat milk and cheese. • Try cooking with herbs, spices, lemon juice instead of butter, margarine or ghee |

MONO UNSATURATED FATS INTAKE AND WEIGHT GAIN TENDENCY

Monounsaturated fatty acids include omega-7 and omega 9 fatty acids. They are associated with anti-inflammatory properties, lowering blood pressure, maintaining triglyceride levels. MUFAs are also found to benefit skin health as they balance water levels and provide ceramides for skin renewal. The Mediterranean diet is rich in MUFA. People of certain genetic type tend to gain weight upon consuming MUFA rich foods than others. These individuals can better maintain weight by balancing the amount of MUFA in their diets. Given that MUFA is beneficial for overall health and particularly heart health, individuals with the weight gain genotype can increase the amount of exercise to compensate for the increased risk of weight gain.



High: Highly likely to gain weight with high MUFA intake

| Genes tested | Recommendations |
|-----------------------------|--|
| <i>NR1D1, ADIPOQ, PPARG</i> | <ul style="list-style-type: none"> • You may have a genetic tendency for higher BMI on a high MUFA diet. • Include MUFA rich food sources in moderation. • Olive oil, avocado, olives, almonds, peanuts are rich in MUFA. • Though a diet rich in MUFA might not help in weight maintenance, it is recommended to include MUFA rich foods in the diet to improve heart health. • Weight gain can be prevented by increasing physical activity |

POLY UNSATURATED FATS INTAKE AND WEIGHT GAIN TENDENCY

Omega 3s are important for brain and heart health as they reduce blood pressure and triglyceride levels. They are also important for skin and vision health. Omega 6s help in bone health and in stimulating hair growth. People of certain genetic type tend to gain weight upon consuming PUFA rich foods than others. These individuals can better maintain weight by lowering the amount of PUFA in their diets or increasing physical activity. Importantly, targeting the Omega3: Omega6 ratio is recommended, with preference towards Omega 3.



Low: Less likely to gain weight with high PUFA intake

| Genes tested | Recommendations |
|--------------|---|
| <i>BDNF</i> | <ul style="list-style-type: none"> You may be less likely to have a higher BMI on a high PUFA diet. Include PUFA-rich food in your diet. Grapeseed oil, canola oil, soybean oil, chia seeds, tuna and mackerel are rich in PUFA. |

PROTEIN INTAKE AND WEIGHT LOSS TENDENCY

Protein is an important building block for bones, skin, blood, cartilage and muscles, and it is present in every cell in the body. Nails, hair, enzymes, hormones and other body chemicals consist of large amounts of protein. Moreover, our body utilises protein to build and repair tissues. People of certain genetic type tend to benefit more in terms of weight maintenance with high protein intake than others



Moderate: Likely to lose weight on high protein intake

| Genes tested | Recommendations |
|--------------|--|
| <i>FTO</i> | <ul style="list-style-type: none"> You may have a genetic tendency to lose moderate weight on a high protein diet. Eat a balanced diet. Include proteins in diet (eggs, chicken breast, tuna, cottage cheese, greek yogurt, almonds, oats, broccoli, quinoa) as they are highly satiating, leading to reduced hunger and appetite |

TENDENCY TO REGAIN WEIGHT

People of certain genetic type tend to quickly regain weight after having been on a weight loss program. These individuals need to continue adherence to exercise and diet program to maintain optimal weight.



Moderate: Likely to regain weight after weight loss

| Genes tested | Recommendations |
|---|--|
| <p><i>PPARG, ADIPOQ, TFAP2B, BDNF</i></p> | <ul style="list-style-type: none"> • You may have a genetic tendency for moderate weight regain after an interventional strategy. • Follow healthy lifestyle and eating pattern to ensure better weight maintenance. • Do not skip breakfast as a good breakfast curbs hunger, avoid processed food and sugar sweetened beverages. • Engage in 30-60 minutes of physical activity. • Maintain a balanced diet and include more of whole grains, nuts, fruits and vegetables to your diet. • Have a food journal and record your daily food habits which will help you to monitor your food consumption |

VITAMIN A NEEDS

Vitamin A is required for clear vision, healthy skin and enhanced immunity. Animal sources provide Vitamin A in the form of retinol, while some plant sources provide the precursor of Vitamin A in the form of carotenenes, which in turn must be converted to retinol. People of certain genetic type need more Vitamin A in their diet due to less efficient conversion of carotenoids to retinol.



Need more: Significantly increase vitamin A intake

| Genes tested | Recommendations |
|-----------------------------|---|
| <p><i>BCMO1, BCMO11</i></p> | <ul style="list-style-type: none"> • You may have a genetic tendency to have low vitamin A levels. • Meet your daily requirements for Vitamin A. • Measure serum Vitamin A level, if below normal even after meeting RDA requirements; consult a physician. • Include carrots, sweet potato, pumpkin, green leafy vegetables, parsley, basil, coriander, milk, fish and bell peppers in daily diet. |

VITAMIN B12 NEEDS

Vitamin B12 is actively involved in red blood cell maturity and its deficiency can lead to pernicious anemia and general fatigue. It also helps in the removal of homocysteine from the cells. People of certain genetic type need more Vitamin B12 in their diet due to lower levels in the body.



Need more: Moderately increase vitamin B12 intake

| Genes tested | Recommendations |
|---------------------------------|---|
| <i>TCN1, CUBN, FUT2, RASIP1</i> | <ul style="list-style-type: none">• You may have a genetic tendency for moderately low vitamin B12 levels.• Meet your daily requirements for Vitamin B12.• Measure serum Vitamin B12 level, if below normal even after meeting RDA requirements; consult a physician.• Vitamin B12 rich foods include fish and seafood.• Also seaweed, eggs, poultry, meat and dairy products provide this nutrient |

VITAMIN B6 NEEDS

Vitamin B6 is required for the proper utilization of sugars, fats and proteins in the body. It also protects the cells against glycation-induced damage. People of certain genetic type need more Vitamin B6 in their diet as they lack the ability to fully metabolize this vitamin leading to its low levels in the body.



Normal: Maintain normal Vitamin B6 intake

| Genes tested | Recommendations |
|---------------------|--|
| <i>NBPF3, ALPL1</i> | <ul style="list-style-type: none">• You may have a genetic tendency for normal vitamin B6 levels.• Meet your daily requirements for Vitamin B6.• Measure serum Vitamin B6 level, if below normal even after meeting RDA requirements; consult a physician.• Vitamin B6 rich foods include whole grain products, nuts and seeds, fish, pork and meat |

VITAMIN B9 NEEDS

Vitamin B9 or folate plays a major role in DNA synthesis and repair. It is also essential for the conversion of homocysteine to methionine. Excess accumulation of homocysteine can be harmful. People of certain genetic type need more Vitamin B9 in their diet due to lower folate levels and an inefficient enzymatic conversion of homocysteine to methionine.



Normal: Maintain normal Vitamin B9 intake

| Genes tested | Recommendations |
|---------------------|---|
| <i>MTHFR</i> | <ul style="list-style-type: none">• You may have a genetic tendency for normal vitamin B9 levels.• Meet your daily requirements for Vitamin B9.• Measure serum Vitamin B9 level, if below normal even after meeting RDA requirements; consult a physician.• Vitamin B9 rich foods include green leafy vegetables, dark coloured fruits (such as oranges, peaches, broccoli, papaya, grapefruit, strawberries, beans, peas, lentils, avocados, okra, sunflower seeds, peanuts, flaxseeds, almonds, cauliflower, corn, celery, carrots and fortified grains) |

VITAMIN C NEEDS

Vitamin C is a potent antioxidant and is essential for enhanced immunity. People of certain genetic type need more Vitamin C in their diet due to lower levels in the body.



Normal: Maintain normal Vitamin C intake

| Genes tested | Recommendations |
|---------------------|--|
| <i>SLC23A1</i> | <ul style="list-style-type: none">• You may have a genetic tendency for normal vitamin C levels.• Meet your daily requirements for Vitamin C.• Measure serum Vitamin C level, if below normal even after meeting RDA requirements; consult a physician.• Vitamin C rich foods include agathi, cabbage, coriander leaves, drumstick leaves, capsicum, guava, green chillies, orange and broccoli |

VITAMIN D NEEDS

Vitamin D is essential for the absorption of calcium from the intestine and also for enhanced immunity. Our body can synthesize sufficient Vitamin D from cholesterol when the skin is exposed to adequate amounts of sunlight. People of certain genetic type need more Vitamin D in their diet due to its inefficient synthesis in our body.



Need more: Moderately increase vitamin D intake

| Genes tested | Recommendations |
|---|--|
| <p><i>GC2, GC1, CYP2R1, CYP27B1, VDR, NADSYN1, GC</i></p> | <ul style="list-style-type: none"> • You may have a genetic tendency for moderately low vitamin D levels. • Meet your daily requirements for Vitamin D. • Include calcium rich foods in the diet to improve absorption of vitamin D. • Measure serum Vitamin D level, if below normal even after meeting RDA requirements; consult a physician. • Calcium rich food sources are chia seeds, beans, lentils, almonds, spinach, tofu, milk & milk products, eggs, mushrooms and finger millets. |

VITAMIN E NEEDS

Vitamin E is an antioxidant and it defends our body against free radical damage and protects polyunsaturated fatty acids from oxidation. People of certain genetic type need more Vitamin E in their diet due to inefficient transport and lower plasma levels of Vitamin E.



Need more: Significantly increase vitamin E intake

| Genes tested | Recommendations |
|--|--|
| <p><i>SCARB1, CD36, CD362, CYP4F2, CD361, ZPR1, TTPA, intergenic</i></p> | <ul style="list-style-type: none"> • You may have a genetic tendency for low vitamin E levels. • Meet your daily requirements for Vitamin E. • Measure serum Vitamin E level, if below normal even after meeting RDA requirements; consult a physician. • Sunflower seeds, olive oil, wheat germ oil, spinach, avocados, almonds, broccoli and shrimps are rich in vitamin E |

VITAMIN K NEEDS

Vitamin K plays an important role in helping blood clotting process and in preventing excessive bleeding. People of certain genetic type need enhanced Vitamin K supplementation to maintain adequate levels in blood.



Need more: Significantly increase vitamin K intake

| Genes tested | Recommendations |
|------------------------------------|---|
| <p><i>GGCX, CYP4F2, VKORC1</i></p> | <ul style="list-style-type: none"> • You may have a genetic tendency for low vitamin K levels. • Meet your daily requirements for vitamin K. • Measure serum vitamin K level, if below normal even after meeting RDA requirements; consult a physician. • Vitamin K rich foods include Brussels sprouts, cabbage, prunes, spring onions and green leafy vegetables. |

CALCIUM NEEDS

Calcium is the most abundant mineral in the body, essential for maintaining the strength and structure of bones and teeth and certain metabolic functions. Both higher and lower calcium levels can have important consequences for health. People of certain genetic type tend to have higher serum calcium levels and can restrict their calcium intake.



Need more: Significantly increase Calcium intake

| Genes tested | Recommendations |
|---|--|
| <p><i>DGKD1, DGKH, GCKR, TTC39B, WDR81, GATA3, DGKD, CYP24A1, CASR, CASR1, CARS</i></p> | <ul style="list-style-type: none"> • You may have a genetic tendency for low calcium levels. • Meet your daily requirements for Calcium (1300mg per day) Measure serum calcium level, if below normal even after meeting RDA requirements; consult a physician. • For adults between 19 and 50 years of age, calcium intake should not exceed 2500mg per day. • For adults older than 50 years, calcium intake should not exceed 2000 mg per day. • Include calcium rich foods such as amaranth leaves, almonds, mustard seeds, sunflower seeds, finger millets, sesame seeds, broccoli and dairy (subject to lactose tolerance recommendation) |

CHOLINE NEEDS

Choline is a macronutrient which plays an important role in liver function, nerve function, normal brain development, muscle movement and in supporting a healthy metabolism. People with a genetic variant in the PEMT gene and other genes are likely to experience adverse health consequences when fed a low choline diet. Hence supplementation is recommended for such individuals.



Need more: Significantly increase Choline intake.

| Genes tested | Recommendations |
|---------------------|---|
| <i>MTHFD1, PEMT</i> | <ul style="list-style-type: none">• You may have a genetic tendency for low choline levels.• Meet your daily requirements for choline.• Measure serum choline level, if below normal even after meeting RDA requirements; consult a physician.• Choline rich foods include eggs, liver, meat, pasta and shellfish. |

COPPER NEEDS

Copper is necessary for the absorption of iron, in the synthesis of haemoglobin and in the maintenance of connective tissue, brain, heart and other organs. People of certain genetic types need more copper.



Need more: Significantly increase Copper intake

| Genes tested | Recommendations |
|------------------------|--|
| <i>SELENBP1, SMIM1</i> | <ul style="list-style-type: none">• You may have a genetic tendency for low copper levels.• Meet your daily requirements for copper.• Measure serum copper level, if below normal even after meeting RDA requirements; consult a physician.• Copper rich foods include sunflower seeds, almonds, dried apricots, dark chocolates and lentils. |

IRON NEEDS

Iron is essential for oxygen transport through the blood. Its deficiency leads to anemia. People of certain genetic type need more iron in their diet as they have reduced ability to absorb iron from the diet.



Need more: Significantly increase Iron intake

| Genes tested | Recommendations |
|---|---|
| <i>TMPRSS6, TFR2, TF, SLC17A1, TMPRSS61</i> | <ul style="list-style-type: none">• You may have a genetic tendency for low iron levels.• Meet your daily requirements for iron.• Men should consume 8 mg per day, women between 19 and 50 years should consume 18 mg per day and women over 50 years should consume 5 mg per day.• Measure serum iron level, if below normal even after meeting RDA requirements; consult a physician.• Iron rich foods include amaranth leaves, spinach, beans, lentils, chickpeas, peas, soybeans, liver, turkey, pumpkin seeds, broccoli, tofu and dark chocolate |

MAGNESIUM NEEDS

Magnesium helps in maintaining normal nerve and muscle function and helps maintain strong bones. It is also important for regulating blood glucose levels and in the production of energy and amino acids.



Need more: Moderately increase Magnesium intake

| Genes tested | Recommendations |
|--|--|
| <i>MDS1, MUC1, SHROOM3, DCDC5, HOXD9, CASR</i> | <ul style="list-style-type: none">• You may have a genetic tendency for moderately low magnesium levels.• Meet your daily requirements for magnesium.• Measure serum magnesium level, if below normal even after meeting RDA requirements; consult a physician.• Magnesium rich foods include dark leafy greens, nuts, fish, whole grains, avocados and yogurt. |

PHOSPHATE NEEDS

Phosphate is necessary for the formation of bones and teeth and is also used as a building block for several important molecules including DNA. People of certain genetic type need more phosphate in their diet as they have decreased phosphate levels in blood.



Need more: Moderately increase Phosphate intake

| Genes tested | Recommendations |
|---------------------|--|
| <i>TKT, CASR</i> | <ul style="list-style-type: none">• You may have a genetic tendency for moderately low phosphate levels.• Meet your daily requirements for Phosphate.• Measure serum phosphate level, if below normal even after meeting RDA requirements; consult a physician.• Phosphate rich foods include pumpkin seeds, brazil nuts, salmons and shellfish |

ZINC NEEDS

Zinc plays an important role in the proper functioning of the immune system, cell division, cell growth and in the breakdown of carbohydrates. Zinc is also important for the senses of taste and smell.



Need more: Moderately increase Zinc intake

| Genes tested | Recommendations |
|-----------------------------|---|
| <i>MT1A, CA1, NBDY, IL6</i> | <ul style="list-style-type: none">• You may have a genetic tendency for moderately low zinc levels.• Meet your daily requirements for zinc.• Measure serum zinc level, if below normal even after meeting RDA requirements; consult a physician.• Zinc rich foods include flax seeds, kidney beans, pumpkin seeds, watermelon seeds and beef |

SELENIUM NEEDS

Selenium helps in the synthesis of antioxidant enzymes and in maintaining a healthy immune system. People of certain genetic type may benefit from selenium supplementation.



Normal: Maintain normal Selenium intake

| Genes tested | Recommendations |
|--------------|---|
| CBS | <ul style="list-style-type: none"> You may have a genetic tendency for normal selenium levels. Meet your daily requirements for selenium. Measure serum selenium level, if below normal even after meeting RDA requirements; consult a physician. Selenium rich foods brazil nuts, yellow fin tuna, turkey, chicken, white button mushrooms and brown rice. |

ANTIOXIDANT NEEDS

Antioxidants play a key role in reducing the ill effects of 'free radicals' and thereby preventing premature aging, tissue damage and the onset of chronic diseases. They are present in many vegetables, fruits, cereals, green tea, etc. People of certain genetic type have lower efficiency to defend themselves against free radical damage and hence require more antioxidants in their diet.



Need more: Moderately increase antioxidants intake

| Genes tested | Recommendations |
|--|---|
| CAT, NAT1, PON1, PON11, SOD2, XRCC1 | <ul style="list-style-type: none"> You may have a genetic tendency to require moderately more antioxidants. Increase intake of foods rich in antioxidants. Optimum level of antioxidants reduces mortality risk. Foods rich in antioxidants are purple, red and blue grapes, blueberries, nuts, green leafy vegetables, sweet potato, carrots, whole grains and beans |

CAFFEINE CONSUMPTION

People of certain genetic type tend to consume more cups of coffee (>625mg of caffeine) a day. Caffeine is a central nervous system stimulant and the most widely consumed psychoactive drug. Increased coffee consumption has been linked to improved health benefits for fast metabolizers of caffeine. Slow metabolizers are prone to increased risk of heart disease with higher caffeine intake.



Moderate: Likely to consume moderate amount of caffeine

Genes tested

Recommendations

AHR, AHR1, CYP1A1, CYP1A2, EFCAB5, MLXIPL, ABCG2, BDNF

- You have a genetic tendency to consume moderate amount of coffee.

CAFFEINE METABOLISM

People of certain genetic type are slow metabolizers of caffeine and may experience symptoms such as palpitations and anxiety upon consuming more than 1 to 2 cups of coffee a day. These individuals may also be at a higher risk of heart disease with increased coffee intake.



Slow: Likely to be a slow metabolizer of caffeine

Genes tested

Recommendations

CYP1A2

- You have a genetic tendency to be a slow metabolizer.
- Restrict coffee intake up to 2 cups a day.
- Choose decaffeinated coffee and other low caffeine beverages.
- May be at higher risk of heart attack when more than 2 cups of coffee are consumed everyday.
- Drink green tea instead.
- Caffeine is present in coffee, energy drinks, in colas and a variety of other foods and beverages.

GLUTEN SENSITIVITY

People of certain genetic type may have lower tolerance to gluten, a protein found in wheat, barley and rye. Some people experience symptoms like abdominal cramps, bloating, "foggy mind", depression, headaches, pain in the bone or joint, diarrhea or constipation and chronic fatigue when they have gluten in their diet but may not test positive for serological determination of celiac disease. When gluten is removed from their diet, these symptoms subside, this condition is known as gluten sensitivity. In some cases (~1% of western population), it may lead to celiac disease.



Gluten Sensitive: Increased risk of being gluten sensitive

| Genes tested | Recommendations |
|--|--|
| <p><i>HLA DQ2.2 (M3), HLA DQ2.2 (M1), HLA DQ2.2 (M2), HLA DQ 7, HLA-DQ 2.5, HLA DQ 8</i></p> | <ul style="list-style-type: none"> • You carry genetic markers associated with increased risk of gluten sensitivity. • However, not all individuals who carry these markers are found to have gluten sensitivity. • Gluten sensitivity needs to be further confirmed by the presence of symptoms. • please consult your physician or dietitian to confirm gluten sensitivity, before starting on a gluten free diet. |

LACTOSE INTOLERANCE

People of certain genetic type stop producing the enzyme lactase in late childhood. Lactase is needed to breakdown the sugar lactose present in milk. These individuals may experience gastrointestinal symptoms upon consuming large quantities of milk as adults.



Intolerant: Likely to be lactose intolerant

| Genes tested | Recommendations |
|--------------------|--|
| <p><i>MCM6</i></p> | <ul style="list-style-type: none"> • You are likely to be lactose intolerant. • Restrict dairy products in the diet. • Limit consumption of whole milk to 1 cup per day and try to distribute the intake throughout the day. • Lactose free milk can be consumed instead of regular milk. • Consume milk along with meals. • Include fermented milk products such as yoghurt that contain lactic acid. • Include non-dairy calcium & vitamin D rich foods in your diet to meet your calcium and vitamin D needs |

SALT INTAKE AND BLOOD PRESSURE SENSITIVITY

Sodium is an essential electrolyte present in the extra cellular fluid. It regulates osmosis and maintains fluid levels within the cell and it also plays an important role in enzyme functions and contraction of muscles. The American Heart Association (AHA) recommends not more than 2300 mg of salt per day; however, most people eat too much salt. On an average 3400 mg of salt is consumed with most of it coming from processed foods. People of certain genetic type will have higher blood pressure in response to high salt consumption. Nearly 50% of hypertensive people are salt-sensitive, which is associated with a rise in BP with salt intake. These individuals will tend to reduce BP with lower salt intake, which is recommended.



Moderate: Likely to have lower BP on low salt diet

| Genes tested | Recommendations |
|--------------------------------------|---|
| <i>NPPA, AGT, ACE, CYP11B2, SGK1</i> | <ul style="list-style-type: none">• You have a genetic tendency to have a slightly above average decrease in blood pressure on a low salt diet.• Moderate to low salt intake recommended |

RIBOFLAVIN AND BLOOD PRESSURE RESPONSE

Riboflavin, also known as vitamin B2, is a water soluble vitamin. Nerves and brain need riboflavin to function properly. It is also required for healthy skin, hair, eyes and liver. people of certain genetic types will have lower blood pressure in response to high riboflavin intake.



Insensitive: Less likely to have lower BP on high Riboflavin intake

| Genes tested | Recommendations |
|---------------------|--|
| <i>MTHFR</i> | <ul style="list-style-type: none">• You have a genetic tendency to have no effect on blood pressure on increased intake of riboflavin.• Though blood pressure levels are not affected by increasing intake of riboflavin, its deficiency can lead to symptoms like burning mouth, angular cheilitis, anemia, and vision problem.• Include foods rich in riboflavin like eggs, liver, dairy products and enriched flour in your diet to ensure optimum level of vitamin B2 in the body. |

ALCOHOL FLUSH

People of certain genetic type may experience symptoms like redness of the face and neck upon consuming alcohol due to reduced clearance of acetaldehyde which is produced in the body upon consuming alcohol. Though alcohol avoidance per limitation is recommended for all, people with the alcohol flush genotype may be at higher health risk upon alcohol consumption



Low: Less likely to experience alcohol flush

| Genes tested | Recommendations |
|---------------------|---|
| <i>ALDH2</i> | <ul style="list-style-type: none">• You have a low genetic tendency to experience alcohol flush.• However, do remember alcohol consumption is a risk factor for many health conditions.• Consume alcohol in moderation. |

الشروط والأحكام وإخلاء المسؤولية



- جميع التحاليل المعلن عنها في متجر بندرجين الإلكتروني ليست تشخيصية وتنحصر الفائدة الإكلينيكية لها مع الأنماط والسمات الصحية لكل شخص بشكل منفصل عن الآخرين. يجب مراجعة الطبيب المختص قبل اتخاذ أي إجراء يتعلق بنتائج هذه التحاليل حيث لن يكون مركز بندرجين الطبي مسؤولاً عن أي إجراء يقوم به العميل استناداً على نتائج هذه التحاليل.
- لا يقوم مركز بندرجين الطبي (فرع شركة مشيخ الطبية أو معهد البحوث والاستشارات بجامعة طيبة) بتقديم أي مشورة طبية بغرض التشخيص ولكن بدلاً عن ذلك فإننا نقوم بتزويدك بمعلومات علمية لفهم وتقييم المخاطر والفوائد الصحية المرتبطة بالنمط الوراثي الخاص بك. مركز بندرجين الطبي يبحث على استشارة الطبيب المختص أو أخصائي التغذية السريرية أو أخصائي الصحة العامة أو الممارس الصحي المؤهل للإجابة على الأسئلة الشخصية الخاصة بك.
- المعلومات التي تم تزويدك بها لا تلغي أو تستبدل أي تشخيص طبي أو نصيحة طبية يمنحها لك الطبيب المختص. كما أن المعلومات العلمية المقدمة لك ليست تشخيصاً وراثياً لأي حالة مرضية. لذلك يجب على الأفراد الذين لديهم مخاوف محددة بشأن حالتهم الصحية أو حالة مرضية محتملة أو أي معلومة خاصة بالاختبارات الجينية التشخيصية أن يقوموا باستشارة الطبيب المختص.
- إن المعلومات المقدمة من مركز بندرجين الطبي ليست شاملة ولا مطلقة، وقد لا تنطبق على حالة الشخص الفردية إذا أخذنا في عين الاعتبار أن المعلومات العلمية في وقت من الأوقات أصبحت غير دقيقة أو قديمة بسبب التقدم العلمي الجديد في مجال التحاليل الجينية والوراثية منذ تاريخ عمل ذلك التحليل. وبمجرد إرسال التقرير لك، فإن مركز بندرجين الطبي لن يقوم بإرسال أي معلومات أو تعليقات أو تحديثات أخرى بشأن التطورات الجديدة التي يتم اكتشافها لاحقاً.
- يُشكّل التقرير المرسل لك كامل نطاق المعلومات التي سيتم تقديمها لك بموجب هذه الاتفاقية. ولا يقدم مركز بندرجين الطبي أي ضمانات أو إقرارات، صريحة أو ضمنية، بأن هذه الخدمة سوف تلي متطلباتك أو توقعاتك حسب نتائج التقرير. كما أن مركز بندرجين الطبي يقوم بإخلاء مسؤوليته عن أي مضاعفات صحية مترتبة على أي إجراء طبي يقوم العميل بإجرائه بناءً على نتائج التقارير المرسلة له.

بعض المراجع العلمية الحديثة في مجال تحليل السمات الجينية



- The integration of epigenetics and genetics in nutrition research for CVD risk factors. Ma Y, Ordovas JM. Proc Nutr Soc. 2017 Aug;76(3):333-346. ([Link](#))
- Genetic Variations Associated with Vitamin A Status and Vitamin A Bioavailability. Borel P, Desmarchelier C. Nutrients. 2017 Mar 8;9(3). ([Link](#))
- Utilizing nutritional genomics to tailor diets for the prevention of cardiovascular disease: a guide for upcoming studies and implementations. Corella D, Coltell O, Mattingley G, Sorlí JV, Ordovas JM. Expert Rev Mol Diagn. 2017 May;17(5):495-513. ([Link](#))
- Gene-Diet Interaction and Precision Nutrition in Obesity. Heianza Y, Qi L. Int J Mol Sci. 2017 Apr 7;18(4). ([Link](#))
- Nutritional modulation of metabolic inflammation. Kirwan AM, Lenighan YM, O'Reilly ME, McGillicuddy FC, Roche HM. Biochem Soc Trans. 2017 Aug 15;45(4):979-985. ([Link](#))
- A multifactorial analysis of obesity as CVD risk factor: Use of neural network based methods in a nutrigenetics context. Valavanis IK, Mougiakakou SG, Grimaldi KA, Nikita KS. BMC Bioinformatics. 2010 Sep 8;11:453. ([Link](#))
- Recent Research in the Genetics of Exercise Training Adaptation. Venezia AC, Roth SM. Med Sport Sci. 2016;61:29-40. ([Link](#))
- Genomic and transcriptomic predictors of response levels to endurance exercise training. Sarzynski MA, Ghosh S, Bouchard C. J Physiol. 2017 May 1;595(9):2931-2939 ([Link](#))
- Advances in Exercise, Fitness, and Performance Genomics in 2015. Sarzynski MA, Loos RJ, Lucia A, Pérusse L, Roth SM, Wolfarth B, Rankinen T, Bouchard C. Med Sci Sports Exerc. 2016 Oct;48(10):1906-16. ([Link](#))